

CONTROVERSY ABOUT QUANTUM TIME DECAY

W.O. Amrein
Département de Physique Théorique
Université de Genève
CH-1211 Genève 4, Switzerland

UGVA-DPT 2001/03-1094

In the recent physics literature there have appeared contradictory statements concerning the behaviour of scattering solutions of the 3-dimensional Schrödinger equation at large times. We clarify the situation and point out that the issue was rigorously resolved in the mathematics literature.

In their papers [1] and [2], García-Calderón, Mateos and Moshinsky considered the nonescape probability $P(t)$ to find a quantum particle, subject to a potential of finite range, still confined in some given finite spatial region at a large time t . They claim that $P(t) \sim t^{-1}$ as $t \rightarrow \infty$. On the other hand Muga, Delgado and Snider in [3] and Cavalcanti in a Comment on [1] (see [4]) obtain a decay of $P(t)$ as t^{-3} . In their Reply [5] to this Comment the authors of [1] reassert that $P(t) \sim t^{-1}$.

The authors cited above seem to be unaware of the fact that, about two decades ago, the question of time decay of solutions of the Schrödinger equation was studied in considerable detail by mathematicians (some relevant references are [6,7] for the 3-dimensional case and [8] for the n -dimensional case, $n = 1, 2, 3, \dots$). The results of these papers clearly imply that $P(t) \sim t^{-3}$ (see e.g. §4 of [6], Theorems 10.3-10.5 in [7] or Theorem 7.6 and Example 7.8 in [8]). More precisely, for a large class of potentials in 1 or 3 dimensions, one has $P(t) = O(t^{-3})$ except possibly for a discrete set of values of the coupling constant (essentially, these exceptional values of the coupling constant are those for which there is a zero energy resonance, and in such a situation the asymptotics of $P(t)$ may be only $O(t^{-1})$). For generic values of the coupling constant the t^{-3} decay of $P(t)$ holds for any bounded spatial domain and any wave packet Ψ such that $\Psi(x, 0)$ (the initial wave function at time $t = 0$) has no bound state components and decays sufficiently rapidly at large $|x|$ (for example, if $n = 3$, such that $\int_{\mathbb{R}^3} (1 + |x|)^{2s} |\Psi(x, 0)|^2 d^3x < \infty$ for some $s > 5/2$). For various initial states Ψ the decay of $P(t)$ will be faster than t^{-3} (several terms in Eq. (7.25) of [8], when applied to Ψ , may be zero).

In the case of s -wave states as considered in [1], the nonescape probability for a ball B_R of radius R (i.e. $B_R = \{x \in \mathbb{R}^3 \mid |x| \equiv r \leq R\}$) is given as follows (see Eqs. (2) and (3) of [1]) :

$$P(t) = \int_0^R \Psi^*(r, t) \Psi(r, t) dr = \int_0^R dr \int_0^R dr' g^*(r, r'; t) \Psi^*(r', 0) \int_0^R dr'' g(r, r''; t) \Psi(r'', 0) ,$$

where $g(r, r'; t)$ denotes the time-dependent retarded Green's function. To calculate $P(t)$ the authors of [1] express the Green's function $g(r, r'; t)$ as an infinite sum of resonant state functions and freely interchange the order of integrations and summations. In the absence of a justification for these interchanges, the expression (14) for $P(t)$ given in [1] (identical with (2.41) in [2]) can not be considered to be exact (in [1] and in the Reply [5], the incorrect time decay $P(t) \sim t^{-1}$ is obtained as a consequence of Eq. (14)). It should also be mentioned that all calculations have to be done for a *fixed* (possibly large) value of

the time parameter t ; the argumentation presented in [5] - invoking an interchange of the long time limit and the integration over the variable r - is beside the point.

The author thanks J.G. Muga for drawing his attention to the disagreement on time decay and for discussions.

References

- [1] G. García-Calderón, J.L. Mateos and M. Moshinsky, Phys. Rev. Lett. **74**, 337 (1995).
- [2] G. García-Calderón, J.L. Mateos and M. Moshinsky, Ann. of Physics **249**, 430 (1996).
- [3] J.G. Muga, V. Delgado and R.F. Snider, Phys. Rev. **B52** 16381 (1995).
- [4] R.M. Cavalcanti, Phys. Rev. Lett. **80**, 4353 (1998).
- [5] G.García-Calderón, J.L.Mateos and M.Moshinsky, Phys. Rev. Lett. **80**, 4354 (1998).
- [6] J. Rauch, Comm. Math. Phys. **61**, 149 (1978).
- [7] A. Jensen and T. Kato, Duke Math. J. **46**, 583 (1979).
- [8] M. Murata, J. Funct. Anal. **49**, 10 (1982).